

## ACUTE TOXICITY SUMMARY

### OZONE

(*triatomic oxygen*)

**CAS Registry Number: 10028-15-6**

#### I. Acute Toxicity Summary (for a 1-hour exposure)

<i>Inhalation reference exposure level</i>	<b>180 <math>\mu\text{g}/\text{m}^3</math></b>
<i>Critical effect(s)</i>	eye irritation and minor changes in lung function tests
<i>Hazard Index target(s)</i>	Eyes; Respiratory System

#### II. Physical and Chemical Properties (HSDB, 1994 except as noted)

<i>Description</i>	colorless to light blue gas
<i>Molecular formula</i>	$\text{O}_3$
<i>Molecular weight</i>	48.0
<i>Density</i>	2.144 g/L @ 0°C (gas)
<i>Boiling point</i>	-111.9°C
<i>Melting point</i>	-192.7°C
<i>Vapor pressure</i>	>760 mm Hg @ 25°C (NIOSH, 1994)
<i>Flashpoint</i>	not applicable
<i>Explosive limits</i>	not applicable
<i>Solubility</i>	insoluble in water; soluble in alkaline solvents, oils
<i>Odor threshold</i>	0.0076-0.036 ppm (15-71 $\mu\text{g}/\text{m}^3$ ) (AIHA, 1989)
<i>Odor description</i>	pungent
<i>Metabolites</i>	unknown
<i>Conversion factor</i>	1 ppm = 1.96 mg/m <sup>3</sup> @ 25°C

#### III. Major Uses or Sources

Ozone is a natural (non-anthropogenic) constituent of the atmosphere with a level between 0.01 and 0.04 ppm. Ozone ( $\text{O}_3$ ) is produced in photochemical reactions of hydrocarbons and nitrogen oxides in the engines of motor vehicles (CARB, 1987) and by certain welding operations. Ozone is used commercially as a disinfectant for air and water. It is also used for bleaching textiles, oils, waxes, and in organic synthesis (ACGIH, 1991).

#### IV. Acute Toxicity to Humans

Impairment of lung function and subsequent impairment of exercise performance were measured in exercising adult athletes (age 19-30) exposed to 0.2 ppm (0.4 mg/m<sup>3</sup>) ozone for 1 hour (Gong *et al.*, 1986). A decrement in post-exercise forced expiratory volume in 1 second (FEV<sub>1</sub>) of

## Determination of Acute Reference Exposure Levels for Airborne Toxicants March 1999

21.6% was observed; a 5.6% decrease in FEV<sub>1</sub> was observed in athletes following a 1-hour exposure to 0.12 ppm (0.24 mg/m<sup>3</sup>) ozone with exercise. Significant reductions in peak minute ventilation, oxygen uptake, and tidal volume were observed in athletes exposed to 0.2 ppm ozone, but not in those exposed to 0.12 ppm.

Healthy young males (age 19-30) exposed to ozone at concentrations as low as 0.12 ppm (0.24 mg/m<sup>3</sup>) for 2.5 hours exhibited statistically significant changes in forced vital capacity (FVC), FEV<sub>1</sub>, forced expiratory flow rates at 75% to 25% of lung volume (FEF<sub>25-75</sub>), and increased coughing (McDonnell *et al.*, 1983). Statistically significant increases in specific airway resistance (SRaw) and reporting of shortness of breath and pain upon deep inspiration were observed in subjects exposed to ozone at concentrations of 0.24 ppm (0.47 mg/m<sup>3</sup>) or greater. A more recent study (McDonnell *et al.*, 1991) reported decrements in FVC, FEV<sub>1</sub>, and significant increases in SRaw and respiratory symptoms in 38 healthy young men following a 6.6-hour exposure to 0.08 ppm (0.2 mg/m<sup>3</sup>) ozone involving 5 hours of exercise.

A statistically significant 3% decrease in FEV<sub>1</sub> was observed in male children (age 8-11) following a 2.5 hour exposure to 0.12 ppm (0.24 mg/m<sup>3</sup>) ozone with intermittent exercise (McDonnell *et al.*, 1985). No significant increase in cough was noted as a result of ozone exposure.

A review by Lippmann (1993) reported that the ozone-associated lower airway response in the normal population engaged in outdoor recreational activity is greatly underestimated by 1 to 2-hour controlled chamber exposure studies, which indicate very little or no functional decrement at 0.120 ppm (249 µg/m<sup>3</sup>) ozone. One study cited by Lippmann (1993) reported significant ozone-associated decrements in FVC, FEV<sub>1</sub>, peak expiratory flow rate (PEFR), FEF<sub>25-75</sub>, and FEV<sub>1</sub>/FVC in healthy adults following outdoor exercise in ambient ozone concentrations of 0.021-0.124 ppm (41-243 mg/m<sup>3</sup>) for an average of 29 minutes (Spektor *et al.*, 1988). In subjects with low ventilation rates (<60 L/minute), the effects observed were about two times greater than those reported in chamber studies using comparable ventilation rates. Recent studies have confirmed that asthmatics react more severely than normal subjects to ozone (Scannell *et al.*, 1996) and that there is a wide variability in spirometric responsiveness (as measured by changes in FVC, FEV<sub>1</sub>, and FEF<sub>25-75</sub>) among individuals to ozone (Weinmann *et al.*, 1995).

### *Predisposing Conditions for Ozone Toxicity*

**Medical:** Persons with preexisting respiratory conditions, such as asthma or chronic obstructive lung disease, may be more sensitive to the adverse effects of ozone exposure (CARB, 1987a). Persons doing vigorous exercise or manual labor outdoors are likely to have increased ventilation rates and to be exposed to a higher dose of ozone and thus may be at increased risk for ozone toxicity.

**Chemical:** Co-exposure to some aeroallergens and respiratory irritants, such as sulfur dioxide, may exacerbate the adverse respiratory effects of ozone in asthmatics (CARB, 1987a).

## **V. Acute Toxicity to Laboratory Animals**

The 3-hour LC<sub>50</sub> values for rats, mice, guinea pigs, and rabbits are reported as 21.8 ppm, 21 ppm, 51.7 ppm, and 36 ppm (42.7, 41, 101, and 71 mg/m<sup>3</sup>) ozone, respectively (Mittler *et al.*, 1956).

A 21% increase in mortality over controls was observed in mice challenged with aerosolized streptococci concurrent with a 3-hour exposure to 0.1 ppm (0.2 mg/m<sup>3</sup>) ozone (Miller *et al.*, 1978). Mice challenged with streptococci immediately following the 3-hour ozone exposure, however, did not exhibit a significant increase in mortality.

Due to the abundance of human exposure studies, additional animal studies were not summarized here.

## **VI. Reproductive or Developmental Toxicity**

No reports of human reproductive or developmental toxicity due to ozone were located in the literature (Shepard, 1994). Increased resorption rates were observed following exposure of pregnant rats to 1.97 ppm (3.86 mg/m<sup>3</sup>) ozone 8 hours per day on days 6-9, 9-12, or 6-15 of gestation (Kavlock *et al.*, 1979). A later study from the same laboratory reported that pregnant rats exposed to 1.0 or 1.5 ppm (2 or 2.9 mg/m<sup>3</sup>) ozone on days 17-20 of gestation had offspring which exhibited retardation of reflex development and slowing in open field behavior (Kavlock *et al.*, 1980).

Veninga (1967) reported blepharophimosis (inability to open the eye to the normal extent) and jaw anomalies in mouse fetuses following maternal exposure to 0.2 ppm (0.4 mg/m<sup>3</sup>) ozone 7 hours per day, 5 days per week. Because the original reference was not available for review, key experimental details (including the days of gestation during which exposure occurred) are not known.

Comparisons of pregnant, lactating, and virgin female rats exposed to 1 ppm (2 mg/m<sup>3</sup>) ozone for 6 hours demonstrated enhanced sensitivity to ozone-induced pulmonary inflammation in pregnant and lactating rats (Gunnison *et al.*, 1992). Pulmonary lavage fluid indicators of inflammation measured include total protein, LDH, total leukocytes, total PMN, and  $\beta$ -glucuronidase activity.

**VII. Derivation of Acute Reference Exposure Level and Other Severity Levels  
(for a 1-hour exposure)**

**Reference Exposure Level (protective against mild adverse effects): 0.09 ppm (180 µg/m<sup>3</sup>)**  
(California Ambient Air Quality Standard)

<i>Study</i>	Gong <i>et al.</i> , 1986; McDonnell <i>et al.</i> , 1983; McDonnell <i>et al.</i> , 1985; California Air Resources Board (CARB), 1987a, 1987b.
<i>Study population</i>	normal adults
<i>Exposure method</i>	inhalation in controlled exposure chambers
<i>Critical effects</i>	decrease in pulmonary function including a 10% decrease in FEV <sub>1</sub>
<i>LOAEL</i>	0.12 ppm (0.24 mg/m <sup>3</sup> ) ozone
<i>NOAEL</i>	not observed
<i>Exposure duration</i>	1 hour
<i>Extrapolated 1 hour concentration</i>	0.12 ppm
<i>LOAEL uncertainty factor</i>	1.3 (margin of safety)
<i>Interspecies uncertainty factor</i>	1
<i>Intraspecies uncertainty factor</i>	1
<i>Cumulative uncertainty factor</i>	1.3 (see below)
<i>Reference Exposure Level</i>	0.09 ppm (0.18 mg/m <sup>3</sup> ; 180 µg/m <sup>3</sup> )

The methodology for developing California Ambient Air Quality Standards differs from that used to develop other acute RELs. The existing CAAQS is based largely upon controlled chamber studies. Inhalation of 0.12 ppm (0.24 mg/m<sup>3</sup>) ozone by normal human subjects in exposure chambers resulted in a decrease in pulmonary function including a 10% decrease in FEV<sub>1</sub>. A margin of safety was added yielding the 1-hour standard of 0.09 ppm (0.18 mg/m<sup>3</sup>). The CAAQS was also designed to protect against eye irritation, a symptom frequently reported when the 1-hour ozone average is 0.1 ppm or greater (although the eye irritation reported may be a result of non-ozone compounds). A recent study (Spektor *et al.*, 1988) reported significant ozone-associated decrements in FVC, FEV<sub>1</sub>, PEFR, FEF<sub>25-75</sub>, and FEV<sub>1</sub>/FVC in healthy adults following outdoor exercise in ambient ozone concentrations of 21-124 ppb (41-243 µg/m<sup>3</sup>) for an average of 29 minutes. In subjects with low ventilation rates (<60 L/minute), the effects observed were about two times greater than those reported in chamber studies using comparable ventilation rates. This new information will be considered when the CAAQS is reevaluated by OEHHA.

**Level Protective Against Severe Adverse Effects**

No recommendation is made due to the limitations of the database.

U.S.EPA (1975) has identified a significant harm level of 0.6 ppm (1.2 mg/m<sup>3</sup>). U.S.EPA states that "at this exposure-time combination [0.6 ppm (1.2 mg/m<sup>3</sup>) ozone for a 1-hour exposure], it is judged that acutely incapacitating symptoms will be experienced by significant portions of the population, especially those engaged in light to moderate exercise, and that the health status of

particularly vulnerable cardiopulmonary subjects may be seriously compromised.” The key study, on which this level is based, is a study of 10 subjects who reported substernal soreness (6/10), cough (8/10), and marked shortness of breath during a 2-hour exposure to 0.75 ppm (1.5 mg/m<sup>3</sup>) ozone involving alternating 15-minute periods of exercise and rest (Bates *et al.*, 1972). The authors concluded that an ozone concentration of 0.75 ppm (1.5 mg/m<sup>3</sup>) produced serious adverse effects under conditions of mild exercise. The choice of the significant harm level is unacceptable as a level protective against severe health effects for exposure of the general public due to the lack of the presentation of a formal protocol for its derivation by U.S.EPA (1975).

### **Level Protective Against Life-threatening Effects**

No recommendation is made due to the limitations of the database.

The NIOSH-IDLH for ozone (NIOSH, 1995) is 10 mg/m<sup>3</sup> (5 ppm) based on acute inhalation toxicity data in humans (Deichmann and Gerarde, 1969; Kleinfeld *et al.*, 1957). According to NIOSH, “Pulmonary edema developed in welders who had a severe acute exposure to an estimated 9 ppm ozone plus other air pollutants (Kleinfeld *et al.*, 1957). It has been reported that on the basis of animal data, exposure at 50 ppm for 60 minutes will probably be fatal to humans (King, 1963).” The derivation of this value is not clearly explained.

### **VIII. References**

(ACGIH) American Conference of Governmental Industrial Hygienists. Documentation of the Threshold Limit Values and Biological Exposure Indices. 6th ed. Cincinnati (OH): ACGIH; 1991. p. 1155-1157.

(AIHA) American Industrial Hygiene Association. Odor thresholds for chemicals with established occupational health standards. Akron (OH): AIHA; 1989. p. 26.

Bates DV, Bell DM, Burnham CD, Hazucha M, Mantha J, Pengelly LD, Silverman F. Short-term effects of ozone on the lung. *J Appl Physiol* 1972;32:176-181. [cited in U.S.EPA, 1975.]

(CARB) California Air Resources Board. Ambient Air Quality Standard for ozone: Health and welfare effects. Staff Report. Sacramento: CARB; September 1987a.

(CARB) California Air Resources Board. Effects of ozone on health. Technical Support Document. Sacramento: CARB; September 1987b.

Deichmann WB, Gerarde HW. Ozone. In: Toxicity of drugs and chemicals. New York (NY): Academic Press, Inc.; 1969. p. 446-448.

Folinsbee LJ, Silverman F, Shepard RJ. Exercise responses following ozone exposure. *J Appl Physiol* 1987;38(6):996-1001.

Determination of Acute Reference Exposure Levels for Airborne Toxicants  
March 1999

Gong H, Bradley PW, Simmons MS, Tashkin DP. Impaired exercise performance and pulmonary function in elite cyclists during low-level ozone exposure in a hot environment. *Am Rev Respir Dis* 1986;134:726-733.

Gunnison AF, Weideman PA, Sobo M. Enhanced inflammatory response to acute ozone exposure in rats during pregnancy and lactation. *Fundam Appl Toxicol* 1992;19:607-612.

(HSDB) Hazardous Substances Data Bank. National Library of Medicine, Bethesda, Maryland (CD-ROM Version). Denver (CO): Micromedex, Inc.; 1994. (Edition expires 7/31/94).

Higgins ITT, D'Arcy JB, Gibbons DI, Avol EL, Gross KB. Effects of exposures to ambient ozone on ventilatory lung function in children. *Am Rev Respir Dis* 1990;141:1136-1146.

Kavlock R, Daston G, Grabowski CT. Studies on the developmental toxicity of ozone. I. Prenatal effects. *Toxicol Appl Pharmacol* 1979;48:19-28.

Kavlock RJ, Meyer E, Grabowski CT. Studies on the developmental toxicity of ozone: postnatal effects. *Toxicol Lett* 1980;5:3-9.

King ME. Toxicity of ozone. V. Factors affecting acute toxicity. *Ind Med Surg* 1963;32:93-94.

Kleinfeld M, Giel C, Tabershaw IR. Health hazards associated with inert-gas-shielded metal arc welding. *AMA Arch Ind Health* 1957;15(1):27-31.

Lippmann M. Health effects of tropospheric ozone: Review of recent research findings and their implications to ambient air quality standards. *J Expo Anal Environ Epidemiol* 1993;3(1):103-129.

McDonnell WF, Horstman DH, Hazucha MJ, Seal E, Haak ED, Salaam SA, *et al.* Pulmonary effects of ozone exposure during exercise: dose-response characteristics. *J Appl Physiol* 1983;54:1345-1352.

McDonnell WF, Chapman RS, Leigh MW, Strobe GL, Collier AM. Respiratory responses of vigorously exercising children to 0.12 ppm ozone exposure. *Am Rev Respir Dis* 1985;132(4):875-879.

McDonnell WF, Kehrl HW, Abdul-Salaam S, Ives PJ, Folinsbee LJ, Devlin RB, *et al.* Respiratory response of humans exposed to low levels of ozone for 6.6 hours. *Arch Environ Health* 1991;46(3):145-150.

McDonnell WF, Muller KE, Bromberg PA, Shy CM. Predictors of individual differences in acute response to ozone exposure. *Am Rev Respir Dis* 1993;147:818-825.

Miller FJ, Illing JW, Gardner DE. Effect of urban ozone levels on laboratory-induced respiratory infections. *Toxicol Lett* 1978;2:163-169.

Determination of Acute Reference Exposure Levels for Airborne Toxicants  
March 1999

Mittler S, Hedrick D, King M, Gaynor A. Toxicity of ozone. I Acute toxicity. *Ind Med Surg* 1956;25:301-306.

(NIOSH) National Institute of Occupational Safety and Health. Chemical listing and documentation of revised IDLH values (as of March 1, 1995). Available at <http://www.cdc.gov/niosh/intridl4.html>.

(NIOSH) National Institute of Occupational Safety and Health Pocket Guide (CD-ROM Version). Denver (CO): Micromedex, Inc.; 1994. (Edition expires 7/31/94).

(NRC) National Research Council. Emergency and continuous exposure limits for selected airborne contaminants. Ozone. Washington, DC: National Academy Press; 1984. p. 99-106.

Scannell C, Chen L, Aris RM, Tager I, Christian D, Ferrando R, *et al*. Greater ozone-induced inflammatory responses in subjects with asthma. *Am J Respir Crit Care Med* 1996;154(1)24-29.

Shepard's catalog of teratogenic agents (CD-ROM Version). Denver (CO): Micromedex, Inc.; 1994. (Edition expires 7/31/94).

Spektor DM, Lippmann M, Thurston GD, Liroy PJ, Stecko J, O'Connor G, *et al*. Effects of ambient ozone on respiratory function in healthy adults exercising outdoors. *Am Rev Respir Dis* 1988;138:821-828.

(U.S.EPA) US Environmental Protection Agency. Prevention of air pollution emergency episodes. Part 51 Chapter I, Title 40 of the Code of Federal Regulations. *Federal Register* 1975;40(162):36333-36335.

Veninga, TS. Toxicity of ozone in comparison with ionizing radiation. *Strahlentherapie* 1967;134:469-477. [cited in Shepard's catalog of teratogenic agents, 1994.]

Weinmann GG, Bowes SM, Gerbase MW, Kimball AW, Frank R. Response to acute ozone exposure in healthy men. Results of a screening procedure. *Am J Respir Crit Care Med* 1995;151(1):33-40.